

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Hu p53 (pS392) Purified, Clone: [FP3.2 FPS392], Monoclonal EXB-11-467-C100

| | |
|--------------------------|---|
| Artikelname | Anti-Hu p53 (pS392) Purified, Clone: [FP3.2 FPS392], Monoclonal |
| Artikelnummer | EXB-11-467-C100 |
| Hersteller Artikelnummer | 11-467-C100 |
| Alternativnummer | EXB-11-467-C100 |
| Hersteller | EXBIO |
| Kategorie | Antikörper |
| Applikation | IHC-P, WB |
| Spezies Reaktivität | Human |
| Immunogen | KLH-conjugated phosphopeptide RHKKLMFKTEGPDS[P]D, corresponding to amino acids 378-393 of human p53. |
| Produktbeschreibung | The tumour suppressor protein p53 is a key element of intracellular anticancer protection. It mediates cell cycle arrest or apoptosis in response to DNA damage or to starvation for pyrimidine nukleotides. It is up-regulated in response to these stres... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/ml |
| Klon-Bezeichnung | [FP3.2 FPS392] |
| Isotyp | Mouse IgG1 |
| Puffer | Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |

| | |
|------------------------|---|
| Lagerung | 2°C to 8°C |
| Target-Kategorie | p53 (pS392) |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (paraffin sections): Standard ABC technique (DAB+), pretreatment: high temperature antigen retrieval (microwave, pressure cooker) in 10 mM citrate buffer pH 6.0 or 1 mM EDTA-NaOH buffer pH 8.0, recommended dilution: 10 µg/ml, incubation: 1 hour at RT, or overnight at 4C, positive tissue: breast carcinoma with high level of wild-type p53. Western blotting: recommended dilution: 1 µg/ml. |