

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Helios Purified, Clone: [22F6], Monoclonal EXB-11-771-C100

Artikelname	Anti-Helios Purified, Clone: [22F6], Monoclonal
Artikelnummer	EXB-11-771-C100
Hersteller Artikelnummer	11-771-C100
Alternativnummer	EXB-11-771-C100
Hersteller	EXBIO
Kategorie	Antikörper
Applikation	FC
Spezies Reaktivität	Human, Mouse
Immunogen	Peptide corresponding to the amino acids 51-107 of Helios
Produktbeschreibung	Helios, also known as IKZF2 (Ikaros family zinc finger protein 2) is a hematopoietic-specific transcription factor involved in the regulation of lymphocyte development, together with other members of this family, such as Aiolos and Ikaros. Helios for...
Klonalität	Monoclonal
Konzentration	1 mg/ml
Klon-Bezeichnung	[22F6]
Isotyp	Armenian Hamster IgG
Puffer	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Lagerung	2°C to 8°C

Target-Kategorie	Helios
Antibody Type	Monoclonal Antibody
Anwendungsbeschreibung	Flow cytometry: Recommended dilution: 1-4 µg/ml., Recommended protocol: 1) Perform staining of cell surface markers (CD25, CD4 etc.) for 20 min. at room temperature in the dark, 100 µl of peripheral blood. 2) Add 3 ml of PBS with 1% BSA, centrifuge at 300g and discard the supernatant. Further steps perform on ice and with ice-cold reagents. 3) Resuspend the cells in 5 ml of cold fixation solution (Miltenyi Biotec) and incubate for 30 min. on ice. 4) Centrifuge for 5 min. at 1000 g, 4C, and discard the supernatant. 5) Resuspend the cells in 5 ml of ice-cold PBS with 1% BSA. 6) Centrifuge for 5 min. at 1000 g, 4C, and discard the supernatant. 7) Resuspend the cells in 5 ml of ice-cold permeabilization solution (Miltenyi Biotec) and incubate 5 min. 8) Centrifuge for 5 min. at 1000 g, 4C, and discard the supernatant. 9) Resuspend the cells in µl of ice-cold permeabilization solution and add 20 µl of FcR blocking solution and incubate for 5 min. (4C, in the dark). 10) Perform intracellular staining of Helios for 30 min. (4C, in the dark) with 10 µl of anti-Helios PE antibody (22F6) in 100 µl. 11) Add 2 ml of ice-cold permeabilization solution and incubate for 5 min. at 4C in the dark. 12) Centrifuge for 5 min. at 1000 g, 4C, and discard the supernatant. 13) Resuspend the cells in 3 ml of ice-cold PBS with 1% BSA. 14) Centrifuge for 5 min. at 1000 g, 4C, and discard the supernatant. 15) Resuspend the cells in 150 µl of ice-cold PBS with 1% BSA and measure on a flow cytometry device with appropriate setting as soon as possible. Keep cold until measuring.