

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Hu CD137 APC, Clone: [4B4-1], Monoclonal EXB-1A-143-T100

|                          |   |
|--------------------------|---|
| Artikelname              | Anti-Hu CD137 APC, Clone: [4B4-1], Monoclonal   |
| Artikelnummer            | EXB-1A-143-T100   |
| Hersteller Artikelnummer | 1A-143-T100   |
| Alternativnummer         | EXB-1A-143-T100   |
| Hersteller               | EXBIO   |
| Kategorie                | Antikörper  |
| Applikation              | FC  |
| Spezies Reaktivität      | Human, Primate  |
| Immunogen                | Recombinant human CD137 ectodomain  |
| Konjugation              | APC   |
| Produktbeschreibung      | CD137, also known as TNFRSF9 or 4-1BB, is an inducible costimulatory molecule expressed mainly on activated T cells. Its ligand, known as 4-1BBL, is expressed on activated macrophages, mature B cells, hematopoietic stem cells, and myeloid progenitor ... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [4B4-1]   |
| Isotyp                   | Mouse IgG1 kappa  |
| Puffer                   | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide   |

|                        |   |
|------------------------|---|
| Lagerung               | 2°C to 8°C  |
| Target-Kategorie       | CD137   |
| Antibody Type          | Monoclonal Antibody   |
| Application Verdünnung | Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests. |
| Anwendungsbeschreibung | Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests. |