

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Hu CD3 APC, Clone: [UCHT1], Monoclonal EXB-1A-514-T100

|                          |   |
|--------------------------|---|
| Artikelname              | Anti-Hu CD3 APC, Clone: [UCHT1], Monoclonal   |
| Artikelnummer            | EXB-1A-514-T100   |
| Hersteller Artikelnummer | 1A-514-T100   |
| Alternativnummer         | EXB-1A-514-T100   |
| Hersteller               | EXBIO   |
| Kategorie                | Antikörper  |
| Applikation              | FC  |
| Spezies Reaktivität      | Human, Primate  |
| Immunogen                | human thymocytes followed by Sezary T cells   |
| Konjugation              | APC   |
| Produktbeschreibung      | CD3 complex is crucial in transducing antigen-recognition signals into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of t... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [UCHT1]   |
| Isotyp                   | Mouse IgG1  |
| Puffer                   | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide   |

|                        |   |
|------------------------|---|
| Lagerung               | 2°C to 8°C  |
| Target-Kategorie       | CD3   |
| Antibody Type          | Monoclonal Antibody   |
| Application Verdünnung | Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.   |
| Anwendungsbeschreibung | Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests. Extracellular and intracellular staining. |