

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Hu CD73 APC-Cy7, Clone: [AD2], APC/Cy7, Monoclonal EXB-T4-675-T100

|                          |   |
|--------------------------|---|
| Artikelname              | Anti-Hu CD73 APC-Cy7, Clone: [AD2], APC/Cy7, Monoclonal   |
| Artikelnummer            | EXB-T4-675-T100   |
| Hersteller Artikelnummer | T4-675-T100   |
| Alternativnummer         | EXB-T4-675-T100   |
| Hersteller               | EXBIO   |
| Kategorie                | Antikörper  |
| Applikation              | FC  |
| Spezies Reaktivität      | Human, Primate  |
| Immunogen                | pre-B leukemia cells  |
| Konjugation              | APC/Cy7   |
| Produktbeschreibung      | CD73 (ecto-5'-nucleotidase) is a 70 kDa glycoprotein anchored to the extracellular leaflet of the plasma membrane by GPI. This ecto-enzyme catalyzes dephosphorylation of AMP to adenosine. CD73 is expressed in various types of cells, such as epithelia... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [AD2]   |
| Isotyp                   | Mouse IgG1 kappa  |
| Puffer                   | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide   |

|                        |  |
|------------------------|--|
| Lagerung               | 2°C to 8°C   |
| Target-Kategorie       | CD73   |
| Antibody Type          | Monoclonal Antibody  |
| Application Verdünnung | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |
| Anwendungsbeschreibung | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |