

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Hu CD16 PE-Cy7, Clone: [3G8], PE/Cy7, Monoclonal EXB-T7-646-T100

| | |
|--------------------------|---|
| Artikelname | Anti-Hu CD16 PE-Cy7, Clone: [3G8], PE/Cy7, Monoclonal |
| Artikelnummer | EXB-T7-646-T100 |
| Hersteller Artikelnummer | T7-646-T100 |
| Alternativnummer | EXB-T7-646-T100 |
| Hersteller | EXBIO |
| Kategorie | Antikörper |
| Applikation | FC |
| Spezies Reaktivität | Human, Primate |
| Immunogen | Human neutrophils |
| Konjugation | PE/Cy7 |
| Produktbeschreibung | CD16 (FcgammaRIII) is a 50-65 kDa glycoprotein serving as a low affinity IgG receptor. Human FcgammaRIII is expressed in two forms - FcgammaRIII-A and -B. FcgammaRIII-A is a transmembrane protein of monocytes, macrophages, NK cells and a subset of T ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [3G8] |
| Isotyp | Mouse IgG1 kappa |
| Puffer | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |

| | |
|------------------------|--|
| Lagerung | 2°C to 8°C |
| Target-Kategorie | CD16 |
| Antibody Type | Monoclonal Antibody |
| Application Verdünnung | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |
| Anwendungsbeschreibung | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |