

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit F(ab)2 anti-Cat IgG (H+L)-HRPO, MinX none DNA-SEC-182653

Artikelname	Rabbit F(ab)2 anti-Cat IgG (H+L)-HRPO, MinX none
Artikelnummer	DNA-SEC-182653
Hersteller Artikelnummer	SEC-182653
Alternativnummer	DNA-SEC-182653
Hersteller	dianova
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Feline
Immunogen	Cat IgG whole molecule
Konjugation	HRPO
Format	F(ab')2
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	F(ab)2 Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and exper...

Klonalität	Polyclonal
Konzentration	10.0 mg/mL
Isotyp	Ig
Puffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product is a F(ab') ₂ fragment of an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Cat IgG and Cat Serum. No reaction was observed against anti-Rabbit IgG F(c) or anti-Pepsin.
Formulierung	Lyophilized
Formel	10 mM NaPO ₄ , 150 mM NaCl, pH 7,2, lyophilisate, Azide/BSA free
Target-Kategorie	Cat
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:10,000
Anwendungsbeschreibung	This product has been assayed against 1.0 ug of Cat IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to 1:100,000 of the reconstitution concentration is suggested for this product. This product has been assayed against 1.0 ug of Cat IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to 1:100,000 of the reconstitution concentration is suggested for this product.