

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Rabbit F(ab)2 anti-Dog IgG (F(ab)2)-FITC, MinX none DNA-SEC-182663

Artikelname	Rabbit F(ab)2 anti-Dog IgG (F(ab)2)-FITC, MinX none
Artikelnummer	DNA-SEC-182663
Hersteller Artikelnummer	SEC-182663
Alternativnummer	DNA-SEC-182663
Hersteller	dianova
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FLISA,FACS,IF
Spezies Reaktivität	Canine
Immunogen	Dog IgG F(ab)2 fragment
Konjugation	FITC
Format	F(ab')2
Spezifität	IgG (F(ab')2)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	F(ab)2 Anti-Dog IgG F(ab)2 Fluorescein Antibody generated in rabbit detects Dog F(ab)2. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and sec...
Klonalität	Polyclonal

Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product is a F(ab') <sub>2</sub> fragment of an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-fluorescein, anti-Rabbit Serum, Dog IgG, Dog IgG F(ab') <sub>2</sub> and Dog Serum. No reaction was observed against Dog IgG F(c), anti-Rabbit IgG F(c) or anti-Pepsin.
Formulierung	Lyophilized
Formel	20 mM K <sub>3</sub> PO <sub>4</sub> , 150 mM NaCl, pH 7.2, lyophilisate, Azide/BSA free
Target-Kategorie	Dog
Antibody Type	Secondary Antibody
Application Verdünnung	FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500-1:2,500, IF Microscopy Dilution: 1:1,000-1:5,000
Anwendungsbeschreibung	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.