

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit IgG anti-Goat IgG (H+L)-unconj., MinX Hu DNA-SEC-182844

| | |
|----------------------------------|---|
| Artikelname | Rabbit IgG anti-Goat IgG (H+L)-unconj., MinX Hu |
| Artikelnummer | DNA-SEC-182844 |
| Hersteller Artikelnummer | SEC-182844 |
| Alternativnummer | DNA-SEC-182844 |
| Hersteller | dianova |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | ELISA,IHC,WB |
| Spezies Reaktivität | Goat |
| Immunogen | Goat IgG whole molecule |
| Konjugation | Unconjugated |
| Format | IgG |
| Spezifität | IgG (H+L) |
| Minimale Kreuzreaktivität (MinX) | Human |
| Produktbeschreibung | Anti-Goat IgG Antibody generated in rabbit detects goat IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi ... |
| Klonalität | Polyclonal |

| | |
|------------------------|---|
| Konzentration | 1.53 mg/mL |
| Isotyp | Ig |
| Puffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Reinheit | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Goat IgG and Goat Serum. No reaction was observed against Human Serum Proteins. |
| Formulierung | Liquid (sterile filtered) |
| Formel | 20 mM K ₃ PO ₄ , 150 mM NaCl, pH 7,2, sterile filtered, 0,01% NaN ₃ |
| Target-Kategorie | Goat |
| Antibody Type | Secondary Antibody |
| Application Verdünnung | ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000 |
| Anwendungsbeschreibung | Anti-Goat IgG antibody has been tested by ELISA and is suitable for western blot and immunohistochemistry, as well as other assays requiring lot-to-lot consistency. |