

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Horse IgG (H+L)-Biotin, MinX none DNA-SEC-182969

| | |
|----------------------------------|---|
| Artikelname | Goat IgG anti-Horse IgG (H+L)-Biotin, MinX none |
| Artikelnummer | DNA-SEC-182969 |
| Hersteller Artikelnummer | SEC-182969 |
| Alternativnummer | DNA-SEC-182969 |
| Hersteller | dianova |
| Wirt | Goat |
| Kategorie | Antikörper |
| Applikation | ELISA,IHC,WB |
| Spezies Reaktivität | Equine |
| Immunogen | Horse IgG whole molecule |
| Konjugation | Biotin |
| Format | IgG |
| Spezifität | IgG (H+L) |
| Minimale Kreuzreaktivität (MinX) | no cross-adsorbtion |
| Produktbeschreibung | Anti-Horse IgG Biotin Antibody generated in goat detects horse IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as... |
| Klonalität | Polyclonal |

| | |
|------------------------|--|
| Konzentration | 2.0 mg/mL |
| Isotyp | Ig |
| Puffer | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Reinheit | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Horse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Horse IgG and Horse Serum. |
| Formulierung | Lyophilized |
| Formel | 10 mM NaPO ₄ , 150 mM NaCl, pH 7,2, lyophilisate, 0,01% NaN ₃ |
| Target-Kategorie | Horse |
| Antibody Type | Secondary Antibody |
| Application Verdünnung | ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000 |
| Anwendungsbeschreibung | This product has been assayed against 1.0 ug of Horse IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to 1:100,000 of the reconstitution concentration is suggested for this product. |