

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Goat IgG anti-Human IgG+IgM+IgA (H+L)-Alk. Phos., MinX none DNA-SEC-182998

Artikelname	Goat IgG anti-Human IgG+IgM+IgA (H+L)-Alk. Phos., MinX none
Artikelnummer	DNA-SEC-182998
Hersteller Artikelnummer	SEC-182998
Alternativnummer	DNA-SEC-182998
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Human
Immunogen	Human IgG, IgA and IgM whole molecule
Konjugation	Alk. Phos.
Format	IgG
Spezifität	IgG+IgM+IgA (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	Anti-Human IgG IgA IgM (H&L) Alkaline Phosphatase Antibody generated in goat detects human (heavy and light chain) immunoglobulin G, A, and M. Immunoglobulin G binds to antigens and can neutralize or opsonize targets, and are predominantly involved i...

Klonalität	Polyclonal
Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine) and anti-Goat Serum. This reagent is suitable for the detection of all human immunoglobulin isotypes, subclasses and chain combinations.
Formulierung	Liquid (sterile filtered)
Formel	50 mM TrisHCl, 150 mM NaCl, 1 mM MgCl, 0,1 mM ZnCl, 50% (v/v) Glycerol, pH 8,0, sterile filtered, 0,1% NaN <sub>3</sub>
Target-Kategorie	Human
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:2,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:3,000
Anwendungsbeschreibung	Anti-Human IgG IgA IgM Alk Phos conjugate is suitable for immunoblotting (western or dot blot), ELISA, immunoelectron microscopy and immunohistochemistry as well as other antibody-based enzymatic assays requiring lot-to-lot consistency.