

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Goat IgG anti-Mouse IgG+IgM+IgA (H+L)-HRPO, MinX none DNA-SEC-183138

Artikelname	Goat IgG anti-Mouse IgG+IgM+IgA (H+L)-HRPO, MinX none
Artikelnummer	DNA-SEC-183138
Hersteller Artikelnummer	SEC-183138
Alternativnummer	DNA-SEC-183138
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Mouse
Immunogen	Anti-Mouse IgG IgA and IgM whole molecule was produced by repeated immunization with Mouse IgG IgA and IgM whole molecule in goat.
Konjugation	HRPO
Format	IgG
Spezifität	IgG+IgM+IgA (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	Anti-Mouse IgG IgA and IgM whole molecule antibody generated in goat detects specifically Mouse IgG IgA and IgM whole molecule. This secondary antibody anti-Mouse is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assay...

Klonalität	Polyclonal
Konzentration	2.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from polyspecific antiserum by immunoaffinity chromatography using Mouse antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase and anti-Goat Serum. This reagent is suitable for the detection of all Mouse immunoglobulin isotypes and chain combinations.
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Thimerosal
Target-Kategorie	Mouse
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:100,000, Immunohistochemistry Dilution: 1:250 - 1:1,500, Western Blot Dilution: 1:500 - 1:3,000
Anwendungsbeschreibung	Suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring lot-to-lot consistency.