

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Mouse IgG (H+L)-HRPO, MinX none DNA-SEC-183170

Artikelname	Goat IgG anti-Mouse IgG (H+L)-HRPO, MinX none
Artikelnummer	DNA-SEC-183170
Hersteller Artikelnummer	SEC-183170
Alternativnummer	DNA-SEC-183170
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Mouse
Immunogen	Anti-Mouse IgG whole molecule was produced by repeated immunization with Mouse IgG whole molecule in goat.
Konjugation	HRPO
Format	IgG
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	Secondary antibodies bind to the primary antibody to assist in detection, sorting and purification of target antigens. To enable detection, the secondary antibody must have specificity for the antibody species and isotype of the primary antibody bein...

Klonalität	Polyclonal
Konzentration	2.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	MOUSE IgG (H&L) Antibody Peroxidase Conjugated was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Mouse IgG and Mouse Serum.
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target-Kategorie	Mouse
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:50,000 - 1:100,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:2,000 - 1:10,000
Anwendungsbeschreibung	Anti-Mouse secondary antibody conjugated to horseradish peroxidase (HRP) generated in goat detects specifically Mouse IgG whole molecule. This anti-Mouse HRP antibody is suitable for ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation, Immunohistochemistry as well as other HRP antibody based assays. Specific conditions for reactivity and signal detection should be optimized by the end user.