

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Rabbit IgG anti-Mouse IgM ( $\mu$ )-Alk. Phos., MinX none DNA-SEC-183284

Artikelname	Rabbit IgG anti-Mouse IgM ( $\mu$ )-Alk. Phos., MinX none
Artikelnummer	DNA-SEC-183284
Hersteller Artikelnummer	SEC-183284
Alternativnummer	DNA-SEC-183284
Hersteller	dianova
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Mouse
Immunogen	Mouse IgM whole molecule
Konjugation	Alk. Phos.
Format	IgG
Spezifität	IgM ( $\mu$ )
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	MOUSE IgM (alpha chain) (RABBIT) Antibody generated in rabbit detects specifically mouse IgM whole molecule. Immunoglobulin M is the largest antibody isotype and the first to be secreted against an initial exposure to antigen. IgM is predominantly pr...
Klonalität	Polyclonal

Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Rabbit Serum, Mouse IgM, and Mouse Serum. No reaction was observed against other mouse heavy or light chain proteins.
Formulierung	Liquid (sterile filtered)
Formel	50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,1% NaN3
Target-Kategorie	Mouse
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:20,000, Immunohistochemistry Dilution: User Optimized, Western Blot Dilution: User Optimized
Anwendungsbeschreibung	Anti-MOUSE IgM (alpha chain) (RABBIT) Antibody is suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring lot-to-lot consistency. Specific conditions for reactivity should be optimized by the end user.