

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat F(ab)2 anti-Human IgG (H+L)-unconj., MinX Bo,Ho,Ms,Rt DNA-SEC-183727

Artikelname	Goat F(ab)2 anti-Human IgG (H+L)-unconj., MinX Bo,Ho,Ms,Rt
Artikelnummer	DNA-SEC-183727
Hersteller Artikelnummer	SEC-183727
Alternativnummer	DNA-SEC-183727
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Human
Immunogen	Anti-Human IgG was produced by repeated immunization with human IgG whole molecule in goat.
Konjugation	Unconjugated
Format	F(ab')2
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	Bovine,Equine,Mouse,Rat
Produktbeschreibung	F(ab)2 Anti-Human IgG (H&L) Antibody generated in goat detects immunoglobulin g from human, both heavy and light chains of the antibody molecule are present. Each IgG has two antigen binding sites. Representing approximately 75% of serum immunoglobul...

Klonalität	Polyclonal
Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	F(ab') ₂ Anti-Human IgG (H&L) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Human IgG and Human Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), or Bovine, Horse, Mouse and Rat Serum Proteins.
Formulierung	Liquid (sterile filtered)
Formel	20 mM K ₃ PO ₄ , 150 mM NaCl, pH 7.2, sterile filtered, 0.01% NaN ₃
Target-Kategorie	Human
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:42,000, Immunohistochemistry Dilution: 1:500 - 1:2,000, Western Blot Dilution: 1:1,000 - 1:5,000
Anwendungsbeschreibung	F(ab) ₂ Anti-Human IgG (H&L) Antibody has been tested by ELISA and dot blot and is suitable for ELISA, immunoblotting, and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring lot-to-lot consistency.