

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat F(ab)2 anti-Mouse IgG+IgM+IgA (H+L)-FITC, MinX Hu, Polyclonal DNA-SEC-183772

Artikelname	Goat F(ab)2 anti-Mouse IgG+IgM+IgA (H+L)-FITC, MinX Hu, Polyclonal
Artikelnummer	DNA-SEC-183772
Hersteller Artikelnummer	SEC-183772
Alternativnummer	DNA-SEC-183772
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	FACS,IF
Spezies Reaktivität	Mouse
Immunogen	Mouse IgG IgA and IgM whole molecule
Konjugation	FITC
Format	F(ab')2
Spezifität	IgG+IgM+IgA (H+L)
Minimale Kreuzreaktivität (MinX)	Human
Produktbeschreibung	F(ab)2 Anti-Mouse IgG IgA IgM (H&L) Fluorescein Antibody generated in goat detects reactivity to Mouse IgG, Mouse IgA, and Mouse IgM subclasses and chains. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing ...

Klonalität	Polyclonal
Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from polyspecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein and anti-Goat Serum. No reaction was observed against anti-Goat IgG F(c), anti-Pepsin or Human Serum Proteins. This product is suitable for the detection of all Human immunoglobulin classes, isotypes and chain combinations.
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target-Kategorie	Mouse
Antibody Type	Secondary Antibody
Application Verdünnung	Flow Cytometry Dilution: User Optimized, IF Microscopy Dilution: 1:500 - 1:2,500
Anwendungsbeschreibung	This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.