

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat F(ab)2 anti-Mouse IgG (Fc)-FITC, MinX Bo,Ho,Hu DNA-SEC-183786

Artikelname	Goat F(ab)2 anti-Mouse IgG (Fc)-FITC, MinX Bo,Ho,Hu
Artikelnummer	DNA-SEC-183786
Hersteller Artikelnummer	SEC-183786
Alternativnummer	DNA-SEC-183786
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	FLISA,FACS,IF
Spezies Reaktivität	Mouse
Immunogen	Mouse IgG F(c) fragment
Konjugation	FITC
Format	F(ab')2
Spezifität	IgG (Fc)
Minimale Kreuzreaktivität (MinX)	Bovine,Equine,Human
Produktbeschreibung	F(ab)2 Anti-Mouse IgG F(c) Fluorescein Antibody was generated in goat and detects specifically Mouse IgG F(c). Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration m...
Klonalität	Polyclonal

Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Goat Serum, Mouse IgG, Mouse IgG F(c) and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Mouse IgG F(ab) or Bovine, Horse and Human Serum Proteins.
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target-Kategorie	Mouse
Antibody Type	Secondary Antibody
Application Verdünnung	FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500 - 1:2,500, Fluorochrome Protein Value: 5.0, IF Microscopy Dilution: 1:1,000 - 1:5,000
Anwendungsbeschreibung	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.