

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Goat F(ab)2 anti-Mouse IgG (F(ab)2)-HRPO, MinX Bo, Ho, Hu, Rb, Rt, Sh DNA-SEC-183792

Artikelname	Goat F(ab)2 anti-Mouse IgG (F(ab)2)-HRPO, MinX Bo, Ho, Hu, Rb, Rt, Sh
Artikelnummer	DNA-SEC-183792
Hersteller Artikelnummer	SEC-183792
Alternativnummer	DNA-SEC-183792
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA, IHC, WB
Spezies Reaktivität	Mouse
Immunogen	F(ab)2 Anti-Mouse IgG F(ab)2 fragment was produced by repeated immunization with Mouse IgG F(ab)2 fragment in goat.
Konjugation	HRPO
Format	F(ab')2
Spezifität	IgG (F(ab')2)
Minimale Kreuzreaktivität (MinX)	Bovine, Equine, Human, Rabbit, Rat, Sheep
Produktbeschreibung	F(ab)2 Anti-Mouse IgG F(ab)2 Peroxidase Antibody generated in goat detects Mouse F(ab)2. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and se...

Klonalität	Polyclonal
Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Mouse IgG, Mouse IgG F(ab') <sub>2</sub> and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Mouse IgG F(c) or Bovine, Horse, Human, Rabbit, Rat and Sheep Serum Proteins.
Formulierung	Lyophilized
Formel	20 mM K <sub>3</sub> PO <sub>4</sub> , 150 mM NaCl, pH 7,2, lyophilisate, 0,01% Gentamicin
Target-Kategorie	Mouse
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:130,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:2,000 to 1:20,000
Anwendungsbeschreibung	F(ab') <sub>2</sub> Anti-Mouse IgG F(ab') <sub>2</sub> fragment is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and signal detection should be optimized by the end user.