

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat F(ab)2 anti-Rabbit IgG (F(ab)2)-HRPO, MinX Bo,Ho,Hu,Ms,Rt,Sh DNA-SEC-183834

Artikelname	Goat F(ab)2 anti-Rabbit IgG (F(ab)2)-HRPO, MinX Bo,Ho,Hu,Ms,Rt,Sh
Artikelnummer	DNA-SEC-183834
Hersteller Artikelnummer	SEC-183834
Alternativnummer	DNA-SEC-183834
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Rabbit
Immunogen	Anti-Rabbit IgG F(ab)2 Antibody was produced in goat by repeated immunization with Rabbit IgG F(ab)2 fragment.
Konjugation	HRPO
Format	F(ab')2
Spezifität	IgG (F(ab')2)
Minimale Kreuzreaktivität (MinX)	Bovine,Equine,Human,Mouse,Rat,Sheep
Produktbeschreibung	F(ab)2 Anti-Rabbit IgG F(ab)2 Peroxidase Antibody generated in goat detects Rabbit F(ab)2. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and ...

Klonalität	Polyclonal
Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	Anti-Rabbit IgG F(ab') ₂ Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Rabbit IgG, Rabbit IgG F(ab') ₂ and Rabbit Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Rabbit IgG F(c) or Bovine, Horse, Human, Mouse, Rat and Sheep Serum Proteins.
Formulierung	Lyophilized
Formel	20 mM K ₃ PO ₄ , 150 mM NaCl, pH 7,2, lyophilisate, 0,01% Gentamicin
Target-Kategorie	Rabbit
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Anwendungsbeschreibung	Suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.