

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

BUBR1 Kinase Antibody, IgG1, Clone: [8G1], Unconjugated, Mouse, Monoclonal Preis auf Anfrage BYT-ORB344429

Article Name	BUBR1 Kinase Antibody, IgG1, Clone: [8G1], Unconjugated, Mouse, Monoclonal Preis auf Anfrage
Biozol Catalog Number	BYT-ORB344429
Supplier Catalog Number	orb344429
Alternative Catalog Number	BYT-ORB344429-100
Manufacturer	Biorbyt
Host	Mouse
Category	Antikörper
Application	ELISA, IF, IP, WB
Species Reactivity	Human
Immunogen	This protein A purified monoclonal antibody was produced by repeated immunizations with a recombinant protein corresponding to amino acid residues 1-350 of human BUBR1 Kinase protein.
Conjugation	Unconjugated
Product Description	BUBR1 Kinase Kinase antibody...
Clonality	Monoclonal
Concentration	1.0 mg/mL
Clone Designation	[8G1]
Isotype	IgG1

NCBI	001202
UniProt	O60566
Buffer	Preservative: 0.01% (w/v) Sodium Azide. Stabilizer: None, Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This Protein A purified antibody is directed against human BUBR1 Kinase protein. The product was purified from tissue culture supernatant by chromatography. This antibody has only been tested on human cells. Reactivity against homologues from other sources is not known.
Form	Liquid (sterile filtered)
Application Dilute	ELISA: 1:5,000 - 1:20,000, IF: 1:200 - 1:1,000, IP: 1:100, WB: 1:200 - 1:1,000
Application Notes	Application Notes: This protein A purified antibody has been tested for use in immunoprecipitation, immunofluorescence staining and western blot and is capable of detecting endogenous protein. Specific conditions for reactivity should be optimized by the end user. Expect a predominant band at ~ 120 kDa corresponding to full-length protein by western blotting in the appropriate cell lysate or extract. Higher MW bands may be seen that may be due to hyperphosphorylation of the protein. The use of HeLa whole cell lysates is recommended as a positive control. For IF microscopy use cells grown on cover slips fixed with 3.5% paraformaldehyde in PBS at pH 6.8. Permeabilize fixed cells with 0.2% Triton X-100 in 25 mM Tris Cl, pH 7.4 containing 0.1% BSA