

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Secretory Component Glycoprotein Antibody / ECM1, Clone: [rECM1/792], Mouse, Monoclonal NSJ-V2484IHC-7ML

Article Name	Recombinant Secretory Component Glycoprotein Antibody / ECM1, Clone: [rECM1/792], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2484IHC-7ML
Supplier Catalog Number	V2484IHC-7ML
Alternative Catalog Number	NSJ-V2484IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Recombinant human protein was used as the immunogen for the recombinant Secretory Component Glycoprotein antibody.
Product Description	This mAb reacts with a reduction-resistant epitope present in both free and IgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free...
Clonality	Monoclonal
Clone Designation	[rECM1/792]
UniProt	Q16610
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	Optimal dilution of the recombinant Secretory Component Glycoprotein antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.