

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Mouse anti Human FAS/APO-1 (CD95), conjugated with PE, IgG1, Clone: [3.22], RPE, Monoclonal NMB-0954

|                            |   |
|----------------------------|---|
| Article Name               | Mouse anti Human FAS/APO-1 (CD95), conjugated with PE, IgG1, Clone: [3.22], RPE, Monoclonal   |
| Biozol Catalog Number      | NMB-0954  |
| Supplier Catalog Number    | 0954  |
| Alternative Catalog Number | NMB-0954  |
| Manufacturer               | NordicMubio   |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | FC  |
| Species Reactivity         | Human   |
| Immunogen                  | Full length CD95 protein  |
| Conjugation                | RPE   |
| Product Description        | Activation of either the 55-kD tumor necrosis factor receptor (TNF-R1) or CD95 (Fas/Apo-1) causes apoptosis of cells and liver failure in mice, and has been associated with human liver disorders. The aim of this study was first to clarify the associa... |
| Clonality                  | Monoclonal  |
| Concentration              | Titered for flow cytometry  |
| Clone Designation          | [3.22]  |

|                   |   |
|-------------------|---|
| Isotype           | IgG1  |
| UniProt           | <a href="#">P25445</a>  |
| Buffer            | Provided as solution in phosphate buffered saline with 0.08% sodium azide   |
| Purity            | Protein A/G Chromatography  |
| Form              | RPE   |
| Formula           | Provided as solution in phosphate buffered saline with 0.08% sodium azide   |
| Application Notes | <p>PBMC: Add 10 µl of MAB/10 6 PBMC in 100 µl PBS. Mix gently and incubate for 15 minutes at 2 to 8C. Wash twice with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze.</p> <p>WHOLE BLOOD: Add 10 µl of MAB/100 µl of whole blood. Mix gently and incubate for 15 minutes at room temperature 20C. Lyse the whole blood. Wash once with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. See instrument manufacturers instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope.</p> |