

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Mouse anti Human CD11b, conjugated with FITC, IgG2a, Clone: [F6.2], Monoclonal NMB-11B2**

Article Name	Mouse anti Human CD11b, conjugated with FITC, IgG2a, Clone: [F6.2], Monoclonal
Biozol Catalog Number	NMB-11B2
Supplier Catalog Number	11B2
Alternative Catalog Number	NMB-11B2
Manufacturer	NordicMubio
Host	Mouse
Category	Antikörper
Application	FC
Species Reactivity	Human
Conjugation	FITC
Product Description	(Mac-1), CD11b/CD18 complex is a leukocyte adhesion receptor to intercellular adhesion molecules (ICAMs). CR3 alpha chain (M.W. 165,000) is on virtually all peripheral blood neutrophils, eosinophils, monocytes and approximately 30% of peripheral bloo...
Clonality	Monoclonal
Concentration	Titered for flow cytometry
Clone Designation	[F6.2]
Isotype	IgG2a

UniProt	<a href="#">P11215</a>
Buffer	Provided as solution in phosphate buffered saline with 0.08% sodium azide and 0.2% carrier protein
Purity	Protein A/G Chromatography
Form	FITC
Formula	Provided as solution in phosphate buffered saline with 0.08% sodium azide and 0.2% carrier protein
Application Notes	<p>PBMC: Add 10 µl of MAB/10<sup>6</sup> PBMC in 100 µl PBS. Mix gently and incubate for 15 minutes at 2 to 8 C. Wash twice with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze.</p> <p>WHOLE BLOOD: Add 10 µl of MAB/100 µl of whole blood. Mix gently and incubate for 15 minutes at room temperature (20C). Lyse the whole blood. Wash once with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. See instrument manufacturers instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope.</p>