

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Mouse anti Human CD3 FITC - CD4 PE, FITC/RPE NMB-B034S

Article Name	Mouse anti Human CD3 FITC - CD4 PE, FITC/RPE
Biozol Catalog Number	NMB-B034S
Supplier Catalog Number	B034S
Alternative Catalog Number	NMB-B034S
Manufacturer	NordicMubio
Host	Mouse
Category	Antikörper
Application	FC
Species Reactivity	Human
Immunogen	CD3=Derived from the hybridization of mouse NS-1 myeloma celss with spleen cells from BALB/c mice immunized with human thymocytes. CD4=Derived from the hybridization of mouse NS-1 myeloma cells with spleen cells from BALB/c mice immunized with human perherial blood T lymphocytes.
Conjugation	FITC/RPE
Product Description	The CD3 epitope is expressed on the epsilon chain of the CD3/T cell antigen receptor (TcR) complex. CD3 is present on 65-85% of thymocytes and has a mitogenic effect on peripheral blood T cells. Identification of human T cells expressing the 22-28,00...
Concentration	Titered for flow cytometry
UniProt	<a href="#">P07766</a>

Buffer	Provided as solution in phosphate buffered saline with 0.08% sodium azide and 0.2% carrier protein
Purity	Protein A/G Chromatography
Form	Bi-Test (FITC/RPE) Reagent
Formula	Provided as solution in phosphate buffered saline with 0.08% sodium azide and 0.2% carrier protein
Application Notes	<p>PBMC: Add 10 µl of MAB/10<sup>6</sup> PBMC in 100 µl PBS. Mix gently and incubate for 15 minutes at 2 to 8°C. Wash twice with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. WHOLE BLOOD: Add 10 µl of MAB/100 µl of whole blood. Mix gently and incubate for 15 minutes at room temperature 20°C. Lyse the whole blood. Wash once with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. See instrument manufacturers instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope.</p>