

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Rabbit anti Human Lysophosphatidic Acid Receptor 3 CT (EDG-7), Unconjugated NMB-X1183P

|                            |  |
|----------------------------|--|
| Article Name               | Rabbit anti Human Lysophosphatidic Acid Receptor 3 CT (EDG-7),<br>Unconjugated   |
| Biozol Catalog Number      | NMB-X1183P   |
| Supplier Catalog Number    | X1183P   |
| Alternative Catalog Number | NMB-X1183P   |
| Manufacturer               | NordicMubio  |
| Host                       | Rabbit   |
| Category                   | Antikörper   |
| Application                | WB   |
| Species Reactivity         | Human  |
| Immunogen                  | Antibody raised against a unique synthetic peptide derived from the<br>C-terminal of the human EDG7 protein.   |
| Conjugation                | Unconjugated   |
| Product Description        | Endothelial Cell Differentiation Gene-7 (EDG-7) belongs to a family of<br>G-protein coupled receptors whose ligands are lysophospholipids.<br>The ligand for EDG-7 is lysophospholipid. There are 8 known<br>members of the EDG receptor family and they are implic... |
| Concentration              | See vial for concentration   |
| Isotype                    | IgG  |
| UniProt                    | <a href="#">Q9UBY5</a>   |

|                   |   |
|-------------------|---|
| Buffer            | Provided as solution in phosphate buffered saline with 0.08% sodium azide   |
| Purity            | Ammonium Sulfate Precipitation  |
| Formula           | Provided as solution in phosphate buffered saline with 0.08% sodium azide   |
| Application Notes | Detects recombinant EDG-7 receptors by Western blot. Due to low expression of EDG receptors, we recommend use of Pierce Femto Signal substrate for western blot development. Optimal concentration should be evaluated by serial dilutions. |