

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit anti Human Sphingosine 1 Phosphate Receptor 5 CT (EDG-8), Unconjugated NMB-X1094P

| Article Name | Rabbit anti Human Sphingosine 1 Phosphate Receptor 5 CT (EDG-8), Unconjugated |
|----------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Biozol Catalog Number | NMB-X1094P |
| Supplier Catalog Number | X1094P |
| Alternative Catalog Number | NMB-X1094P |
| Manufacturer | NordicMubio |
| Host | Rabbit |
| Category | Antikörper |
| Application | WB |
| Species Reactivity | Human |
| Conjugation | Unconjugated |
| Product Description | Endothelial Cell Differentiation Gene-8 (EDG-8) belongs to a family of G-protein coupled receptors whose ligands are lysophospholipids. The ligand for EDG-8 is sphingosine-1-phosphate. There are 8 known members of the EDG receptor family and they are |
| Concentration | See vial for concentration |
| Isotype | IgG |
| UniProt | Q9H228 |
| Buffer | Provided as solution in phosphate buffered saline with 0.08% sodium azide |

| Source | Rabbits were immunized with a unique synthetic peptide derived from the C-terminus of the human EDG-8 protein |
|-------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Purity | Ammonium Sulfate Precipitation |
| Application Notes | Detects recombinant EDG8 receptors by Western blot (5-10 μ g/ml) at 42 kDa. Due to low expression of EDG receptors, we recommend use of Pierce Femto Signal substrate for western blot development. Optimal concentration should be evaluated by serial dilutio |