

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Chicken IgY anti-Mouse IgG (F(ab')2)-FITC, MinX none NMB-AMIGGFAB-F

|                            |  |
|----------------------------|--|
| Article Name               | Chicken IgY anti-Mouse IgG (F(ab')2)-FITC, MinX none                     |
| Biozol Catalog Number      | NMB-AMIGGFAB-F   |
| Supplier Catalog Number    | AM IgG Fab-F   |
| Alternative Catalog Number | NMB-AMIGGFAB-F   |
| Manufacturer               | NordicMubio  |
| Host                       | Gallus   |
| Category                   | Antikörper   |
| Application                | IHC  |
| Species Reactivity         | Mouse  |
| Conjugation                | FITC   |
| Format                     | IgY  |
| Target Specificity         | IgG (F(ab')2)  |
| Cross-Adsorption (MinX)    | no cross-adsorbtion  |
| Product Description        | Chicken anti Mouse IgG (Fab)2, conjugated with FITC, Gallus 0,5 mg...    |
| Concentration              | See vial for concentration.  |
| Buffer                     | Phosphate buffered saline, pH 7.2, 50% glycerol with 0.075% sodium azide |

|                   |   |
|-------------------|---|
| Source            | Chickens were immunized with highly purified normal mouse IgG. After multiple immunizations in Freunds adjuvant, eggs were collected and the IgY fraction prepared. The anti-(Fab)2 fraction was immunoaffinity- purified and conjugated to fluorescein isothio |
| Purity            | Egg-yolk derived IgY Affinity Purified  |
| Form              | Fluorescein conjugated  |
| Formula           | Phosphate buffered saline, pH 7.2, 50% glycerol with 0.075% sodium azide  |
| Antibody Type     | Secondary Antibody  |
| Application Notes | This antibody is suitable for immunohistochemical applications. Optimal working dilutions should be determined for your particular assay conditions. The F/P ratio is between 3 and 6.  |