

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Goat IgG anti-Bovine IgG (H+L)-FITC, MinX Hu,Sw NMB-GAB/IGG(H+L)/FITC

Article Name	Goat IgG anti-Bovine IgG (H+L)-FITC, MinX Hu,Sw
Biozol Catalog Number	NMB-GAB/IGG(H+L)/FITC
Supplier Catalog Number	GAB/IgG(H+L)/FITC
Alternative Catalog Number	NMB-GAB/IGG(H+L)/FITC
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Bovine
Conjugation	FITC
Format	IgG
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	Human,Porcine
Product Description	The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes some reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in bovine serum, as tested by imm
Clonality	Polyclonal
Clone Designation	[Polyclonal]

Buffer	FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added. It is reconstituted by adding 2 ml sterile distilled water, spun down to remove insoluble particles, divided into sm
Source	Purified normal IgG isolated from pooled bovine serum. Freunds complete adjuvant is used in the first step of the immunization procedure.
Formula	FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added.
Antibody Type	Secondary Antibody
Application Notes	ELISA,Immunocytochemistry,Immunohistochemistry (frozen),(In)direct immunofluorescence.