

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Dog IgM (μ)-unconj., MinX none NMB-GAD/IGM(FC)/7S

Article Name	Goat IgG anti-Dog IgM (μ)-unconj., MinX none
Biozol Catalog Number	NMB-GAD/IGM(FC)/7S
Supplier Catalog Number	GAD/IgM(Fc)/7S
Alternative Catalog Number	NMB-GAD/IGM(FC)/7S
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Canine
Conjugation	Unconjugated
Format	IgG
Target Specificity	IgM (μ)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is directed to the Fc subunit of the IgM molecule which expresses strict isotypic (class) specificity. It does not react with IgG, IgA and IgG/Fab or any non-Ig protein in dog serum, as tested by immunoelectrophoresis
Clonality	Polyclonal
Clone Designation	[Polyclonal]

Buffer	Purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove i
Source	Purified IgM isolated from dog serum. Freunds complete adjuvant is used in the first step of the immunization procedure.
Formula	Purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.
Antibody Type	Secondary Antibody
Application Notes	Indirect immunofluorescence, ELISA, Dot blot, Immunoblotting.