

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Goat IgG anti-Dog IgM (μ)-Biotin, MinX none NMB-GAD/IGM(FC)/BIO

A	
Article Name	Goat IgG anti-Dog IgM (μ)-Biotin, MinX none
Biozol Catalog Number	NMB-GAD/IGM(FC)/BIO
Supplier Catalog Number	GAD/IgM(Fc)/Bio
Alternative Catalog Number	NMB-GAD/IGM(FC)/BIO
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Canine
Conjugation	Biotin
Format	IgG
Target Specificity	IgM (μ)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is directed to the Fc subunit of the IgM molecule which expresses strict isotypic (class) specificity. It does not react with any non-Ig protein in dog serum, as tested by immunoelectrophoresis and double radial immuno
Clonality	Polyclonal
Clone Designation	[Polyclonal]

Buffer	Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun d
Source	Purified normal IgM isolated from pooled dog serum. Freunds complete adjuvant is used in the first step of the immunization procedure.
Formula	Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.
Antibody Type	Secondary Antibody
Application Notes	ELISA,Immunocytochemistry,Immunohistochemistry (paraffin),Dot blot,Immunoblotting.