

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat IgG anti-Dog IgM ( $\mu$ )-FITC, MinX none NMB-GAD/IGM(FC)/FITC

Article Name	Goat IgG anti-Dog IgM ( $\mu$ )-FITC, MinX none
Biozol Catalog Number	NMB-GAD/IGM(FC)/FITC
Supplier Catalog Number	GAD/IgM(Fc)/FITC
Alternative Catalog Number	NMB-GAD/IGM(FC)/FITC
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Canine
Conjugation	FITC
Format	IgG
Target Specificity	IgM ( $\mu$ )
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is directed to the Fc subunit of the IgM molecule, which expresses strict (class) specificity. In immunoelectrophoresis and radial in immunodiffusion using various antiserum concentrations against dog serum, a single p...
Clonality	Polyclonal
Clone Designation	[Polyclonal]

Buffer	FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun down
Source	Purified IgM isolated from dog serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formula	FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.
Antibody Type	Secondary Antibody
Application Notes	ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence.