

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Duck albumin, Clone: [Polyclonal], Monoclonal NMB-GADU/ALB

| | |
|----------------------------|---|
| Article Name | Goat anti Duck albumin, Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-GADU/ALB |
| Supplier Catalog Number | GADu/Alb |
| Alternative Catalog Number | NMB-GADU/ALB |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Product Description | The defined antibody specificity is directed to albumin as tested against duck sera. In immunoelectrophoresis and double radial immunodiffusion (Ouchterlony), using various antiserum concentrations against appropriate concentrations of the immunogen,... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those in pooled goat serum. No foreign proteins added. Reconstitute the lyophilized antiserum |
| Source | Highly purified albumin isolated from duck serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |

| | |
|-------------------|--|
| Formula | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those in pooled goat serum. No foreign proteins added. |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 µl serum, plasma or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and serially diluted) in 2 mm diameter peripheral wells. In single radial immunodiffusion and electroimmunodiffusion use 0.5 to 1.0 percent antiserum in the agar gel. |