

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Goat Antiserum anti-Bird IgG+IgM+IgA (H+L)-unconj., MinX none NMB-GADU/IG

Article Name	Goat Antiserum anti-Bird IgG+IgM+IgA (H+L)-unconj., MinX none
Biozol Catalog Number	NMB-GADU/IG
Supplier Catalog Number	GADu/Ig
Alternative Catalog Number	NMB-GADU/IG
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Aves
Conjugation	Unconjugated
Format	Antiserum
Target Specificity	IgG+IgM+IgA (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is directed to the Fc and Fab subunits of the major immunoglobulins. It does not react with any non-lg protein in duck serum, as tested by immunoelectrophoresis and double radial immunodiffusion. In precipitating techn
Clonality	Polyclonal
Clone Designation	[Polyclonal]

Buffer	Delipidated, heat inactivated, lyophilized stable whole serum No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat ser
Source	Purified normal Ig fractions isolated from pooled duck serum.  Freunds complete adjuvant is used in the first step of the immunization procedure.
Formula	Delipidated, heat inactivated, lyophilized stable whole serum No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat ser
Antibody Type	Secondary Antibody
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 $\mu$ l or equivalent against 120 $\mu$ l antiserum. In double radial immunodiffusion use a rosette arrangement with 10 $\mu$ l antiserum in a 3 mm diameter centre well and 2 $\mu$ l serum samples (neat and diluted) in 2