

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Goat Antiserum anti-Guinea Pig IgG (F(ab')2)-unconj., MinX none NMB-GAGP/FAB

Article Name	Goat Antiserum anti-Guinea Pig IgG (F(ab')2)-unconj., MinX none
Biozol Catalog Number	NMB-GAGP/FAB
Supplier Catalog Number	GAGp/Fab
Alternative Catalog Number	NMB-GAGP/FAB
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Guinea pig
Conjugation	Unconjugated
Format	Antiserum
Target Specificity	IgG (F(ab')2)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The antiserum is reacting with the Fab subunits IgG, IgA, IgM and other Ig classes of both light chain type, with their Fab or F(ab)2 subunits and with free light chains of kappa and lambda type. It does not react with any non-Ig protein in guinea pi
Clonality	Polyclonal
Clone Designation	[Polyclonal]

Buffer	Delipidated, heat inactivated, lyophilized stable whole serum No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat ser
Source	Purified Fab of normal IgG isolated from pooled guinea pig serum. Freunds complete adjuvant is used in the first step of the immunization procedure.
Formula	Delipidated, heat inactivated, lyophilized stable whole serum No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat ser
Antibody Type	Secondary Antibody
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 $\mu$ l or equivalent against 120 $\mu$ l antiserum. In double radial immunodiffusion use a rosette arrangement with 10 $\mu$ l antiserum in a 3 mm diameter centre well and 2 $\mu$ l serum samples (neat and diluted) in 2