

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Guinea Pig IgG (F(ab')₂)-FITC, MinX none NMB-GAGP/FAB/FITC

| | |
|----------------------------|---|
| Article Name | Goat IgG anti-Guinea Pig IgG (F(ab') ₂)-FITC, MinX none |
| Biozol Catalog Number | NMB-GAGP/FAB/FITC |
| Supplier Catalog Number | GAGp/Fab/FITC |
| Alternative Catalog Number | NMB-GAGP/FAB/FITC |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Species Reactivity | Guinea pig |
| Conjugation | FITC |
| Format | IgG |
| Target Specificity | IgG (F(ab') ₂) |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | The reactivity of the antiserum is directed to the Fab subunits of intact IgG, IgA IgM and other Ig classes of both light chain types, with their Fab or F(ab) ₂ subunits, and with free light chains of kappa and lambda type. In immunoelectrophoresis an... |
| Clonality | Polyclonal |
| Clone Designation | [Polyclonal] |

| | |
|-------------------|--|
| Buffer | FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun down |
| Source | Purified normal Fab of polyclonal guinea pig IgG. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Form | FITC |
| Formula | FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added as it may interfere with the antibody activity. |
| Antibody Type | Secondary Antibody |
| Application Notes | ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence. |