

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat Antiserum anti-Horse IgG (Fc)-unconj., MinX none NMB-GAHO/IGG(FC)

Article Name	Goat Antiserum anti-Horse IgG (Fc)-unconj., MinX none
Biozol Catalog Number	NMB-GAHO/IGG(FC)
Supplier Catalog Number	GAHo/IgG(Fc)
Alternative Catalog Number	NMB-GAHO/IGG(FC)
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Application	IP
Species Reactivity	Equine
Conjugation	Unconjugated
Format	Antiserum
Target Specificity	IgG (Fc)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is restricted to the Fc part of the IgG molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal horse serum a single precipitin line is obtained which shows ...
Clonality	Polyclonal
Clone Designation	[Polyclonal]

Buffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized
Source	Highly purified normal IgG isolated from pooled horse serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Antibody Type	Secondary Antibody
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 $\mu$ l serum or equivalent against 120 $\mu$ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 $\mu$ l antiserum in 3 mm diameter center well and 2 $\mu$ l serum samples (neat