

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Human alpha-1 lipoprotein, Clone: [Polyclonal], Monoclonal NMB-GAHU/A1LP(HDL)

| Article Name | Goat anti Human alpha-1 lipoprotein, Clone: [Polyclonal], Monoclonal |
|----------------------------|--|
| Biozol Catalog Number | NMB-GAHU/A1LP(HDL) |
| Supplier Catalog Number | GAHu/A1Lp(HDL) |
| Alternative Catalog Number | NMB-GAHU/A1LP(HDL) |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Product Description | The reactivity of the antiserum is restricted to alpha-1 lipoprotein. In immunoelectrophoresis and radial immunodiffusion (Ouchterlony), using various antiserum concentrations against fresh normal human plasma a single precipitin line is obtained whi |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. |
| Source | Alpha-1 lipoproteins are completely different from beta-lipoproteins. In fresh serum only one alpha-1 lipoprotein exists immunologically, but after centrifugation, storage, freezing and thawing two types of alpha-1 lipoprotein with difference in mobility |

| Formula | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. |
|-------------------|---|
| Application Notes | Precipitation assays. The lyophilized antiserum is shipped at ambient temperature and may be stored at +4C, prolonged storage at or below -20C. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Dilutions may be prepared by |