

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat anti Human alpha-2 antiplasmin, Clone: [Polyclonal], Monoclonal NMB-GAHU/APL

Article Name	Goat anti Human alpha-2 antiplasmin, Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-GAHU/APL
Supplier Catalog Number	GAHu/Apl
Alternative Catalog Number	NMB-GAHU/APL
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Product Description	The defined antibody reactivity is restricted to alpha-2 antiplasmin as tested at the level of sensitivity of immuno-precipitation techniques. A single precipitin line is obtained in bidimensional electrophoresis, immunoelectrophoresis and double rad...
Clonality	Monoclonal
Clone Designation	[Polyclonal]
UniProt	<a href="#">P08697</a>
Buffer	Delipidated, heat inactivated, lyophilized, stable whole serum. Sodium azide 1 mg/ml. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized antise
Source	Apha-2 antiplasmin is a single-chain glycoprotein with a molecular weight of about 65,000. The normal concentration in human plasma is 60 µg/ml. Platelets contain significant amounts of the protein, which is released along with platelet factor 4, following

Formula	Delipidated, heat inactivated, lyophilized, stable whole serum. Sodium azide 1 mg/ml. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added.
Application Notes	Precipitation assays. In electroimmunodiffusion the amount of antiserum required in the agarose gel is usually between 1.5 and 2% depending on the test arrangement.