

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Human C3c, conjugated with Biotin, Clone: [Polyclonal], Monoclonal NMB-GAHU/C3C/BIO

| | |
|----------------------------|--|
| Article Name | Goat anti Human C3c, conjugated with Biotin, Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-GAHU/C3C/BIO |
| Supplier Catalog Number | GAHu/C3c/Bio |
| Alternative Catalog Number | NMB-GAHU/C3C/BIO |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Conjugation | Biotin |
| Product Description | In immunoelectrophoresis against fresh human serum, a single precipitin line is obtained in the beta-1 region representing native C3. Against serum containing partly activated C3, a precipitin line is obtained which extends from the beta-1 into the a... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| UniProt | P01024 |
| Buffer | Biotin-coupled purified hyperimmune IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2) No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun down to |

| | |
|-------------------|---|
| Source | C3c is the major fragment resulting from C3 cleavage by C3 convertase and factor I. It is composed of an intact beta chain bound to two fragments of the alpha chain. C3c is isolated and purified from pooled normal human serum. Freund's complete adjuvant |
| Formula | Biotin-coupled purified hyperimmune IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2) No preservative added, as it may interfere with the antibody activity. |
| Application Notes | ELISA, Immunocytochemistry, Immunohistochemistry (paraffin), Dot blot, Immunoblotting. |