

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Human IgA (Fc specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal NMB-GAHU/IGA(FC)/TRITC

Article Name	Goat anti Human IgA (Fc specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-GAHU/IGA(FC)/TRITC
Supplier Catalog Number	GAHu/lgA(Fc)/TRITC
Alternative Catalog Number	NMB-GAHU/IGA(FC)/TRITC
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Conjugation	TRITC
Product Description	The reactivity of the antiserum is directed to the Fc subunit of the IgA molecule which expresses strict isotypic (class) specificity. It does not react with any non-Ig protein in human serum, as tested by immunoelectrophoresis and double radial immu
Clonality	Monoclonal
Clone Designation	[Polyclonal]
Buffer	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun do

Source	Purified polyclonal IgA and monoclonal IgA2 isolated from pooled human serum. Freunds complete adjuvant is used in the first step of the immunization procedure.
Formula	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.
Application Notes	ELISA,Immunocytochemistry,Immunohistochemistry (frozen),(In)direct immunofluorescence.