

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Human IgD (Fc specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal NMB-GAHU/IGD(FC)/TRITC

| | |
|----------------------------|---|
| Article Name | Goat anti Human IgD (Fc specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-GAHU/IGD(FC)/TRITC |
| Supplier Catalog Number | GAHu/IgD(Fc)/TRITC |
| Alternative Catalog Number | NMB-GAHU/IGD(FC)/TRITC |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Species Reactivity | Human |
| Conjugation | TRITC |
| Format | IgG |
| Target Specificity | IgD |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | The reactivity of the antiserum is directed to the Fc subunit of the IgD molecule which expresses strict isotypic (class) specificity. It does not react with any non-Ig protein in human serum, as tested by immunoelectrophoresis and double radial immu... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |

| | |
|-------------------|--|
| Buffer | TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun down |
| Source | Purified monoclonal IgD isolated from pooled human serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Form | TRITC |
| Formula | TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. |
| Antibody Type | Secondary Antibody |
| Application Notes | ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence. |