

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Human IgG (F(ab')2)-TRITC, MinX none NMB-GAHU/FAB/TRITC

Article Name	Goat IgG anti-Human IgG (F(ab')2)-TRITC, MinX none
Biozol Catalog Number	NMB-GAHU/FAB/TRITC
Supplier Catalog Number	GAHu/Fab/TRITC
Alternative Catalog Number	NMB-GAHU/FAB/TRITC
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Human
Conjugation	TRITC
Format	IgG
Target Specificity	IgG (F(ab')2)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is directed to the Fab subunits of intact IgG, IgA IgM and other Ig classes of both light chain types, with their Fab or F(ab)2 subunits, and with free light chains of kappa and lambda type. In immunoelectrophoresis an
Clonality	Polyclonal
Clone Designation	[Polyclonal]

Buffer	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added as it may interfere with he antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun down
Source	Purified Fab of normal IgG isolated form human serum. Freunds complete adjuvant is used in the first step of the immunization procedure.
Formula	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added as it may interfere with he antibody activity.
Antibody Type	Secondary Antibody
Application Notes	ELISA,Immunocytochemistry,Immunohistochemistry (frozen),(In)direct immunofluorescence.