

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat Antiserum anti-Human IgG+IgM+IgA (Fc)-unconj., MinX none NMB-GAHU/IG(FC)

| | |
|----------------------------|---|
| Article Name | Goat Antiserum anti-Human IgG+IgM+IgA (Fc)-unconj., MinX none |
| Biozol Catalog Number | NMB-GAHU/IG(FC) |
| Supplier Catalog Number | GAHu/Ig(Fc) |
| Alternative Catalog Number | NMB-GAHU/IG(FC) |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Species Reactivity | Human |
| Conjugation | Unconjugated |
| Format | Antiserum |
| Target Specificity | IgG+IgM+IgA (Fc) |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | The reactivity of the antiserum is directed to the Fc subunits of the major immunoglobulins. It does not react with any non-Ig protein in human serum, as tested by immunoelectrophoresis and double radial immunodiffusion. In precipitating techniques a... |
| Clonality | Polyclonal |
| Clone Designation | [Polyclonal] |

| | |
|-------------------|--|
| Buffer | Delipidated, heat inactivated, lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum. |
| Source | Purified normal IgG, IgA and IgM fractions isolated from pooled human serum. Freunds complete adjuvant is used in the first step of the immunization procedure. |
| Formula | Delipidated, heat inactivated, lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum. |
| Antibody Type | Secondary Antibody |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 |