

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Goat IgG anti-Human IgG+IgM+IgA (Fc)-TRITC, MinX none NMB-GAHU/IG(FC)/TRITC

Article Name	Goat IgG anti-Human IgG+IgM+IgA (Fc)-TRITC, MinX none
	Code igo dila Haman igo Figir Figir (Fo) Harro, Minor Hone
Biozol Catalog Number	NMB-GAHU/IG(FC)/TRITC
Supplier Catalog Number	GAHu/lg(Fc)/TRITC
Alternative Catalog Number	NMB-GAHU/IG(FC)/TRITC
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Human
Conjugation	TRITC
Format	IgG
Target Specificity	IgG+IgM+IgA (Fc)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is directed to the Fc subunits of the major isotypes of the human immunoglobulin system. No reaction is obtained with purified IgG/Fab or any non-Ig protein of human serum, as tested by immunoelectrophoresis and double
Clonality	Polyclonal
Clone Designation	[Polyclonal]

Buffer	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water.
Source	Purified immunoglobulin fractions containing IgG, IgA and IgM isolated from pooled human serum. Freunds complete adjuvant is used in the first step of the immunization procedure.
Formula	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.
Antibody Type	Secondary Antibody
Application Notes	ELISA,Immunocytochemistry,Immunohistochemistry (frozen),(In)direct immunofluorescence.