

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat Antiserum anti-Human IgG+IgM+IgA+IgD (H+L)-unconj., MinX none NMB-GAHU/IG(GAMD)

Article Name	Goat Antiserum anti-Human IgG+IgM+IgA+IgD (H+L)-unconj., MinX none
Biozol Catalog Number	NMB-GAHU/IG(GAMD)
Supplier Catalog Number	GAHu/Ig(GAMD)
Alternative Catalog Number	NMB-GAHU/IG(GAMD)
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Human
Conjugation	Unconjugated
Format	Antiserum
Target Specificity	IgG+IgM+IgA+IgD (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is directed to the major isotypes of the human immunoglobulin system (classes and both light chain types) including antibodies to common determinants, to class and to the surface determinants of the common Fab portion,...
Clonality	Polyclonal
Clone Designation	[Polyclonal]

Buffer	Delipidated, heat inactivated, lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum.
Source	Purified normal IgG and fractions containing IgA, IgM and homogenous IgD isolated from human serum. Freunds complete adjuvant is used in the first step of the immunization procedure.
Formula	Delipidated, heat inactivated, lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum.
Antibody Type	Secondary Antibody
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2