

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat F(ab')2 anti-Human IgM (μ)-TRITC, MinX none NMB-FGAHU/IGM(FC)/TRITC

Article Name	Goat F(ab')2 anti-Human IgM (μ)-TRITC, MinX none
Biozol Catalog Number	NMB-FGAHU/IGM(FC)/TRITC
Supplier Catalog Number	FGAHu/IgM(Fc)/TRITC
Alternative Catalog Number	NMB-FGAHU/IGM(FC)/TRITC
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Human
Conjugation	TRITC
Format	F(ab')2
Target Specificity	IgM (μ)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The antiserum is reacting with to the Fc subunits of IgM. It does not react with other immunoglobulins or any non-Ig protein in human serum, as tested by immunoelectrophoresis and double radial immunodiffusion. Direct staining of fixed cell and tissu
Clonality	Polyclonal
Clone Designation	[Polyclonal]

Buffer	TRITC-coupled goat F(ab)2 antibodies lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small a
Source	Purified IgM prepared from pooled human serum. Freunds complete adjuvant is used in the first step of the immunization procedure.
Formula	TRITC-coupled goat F(abO)2 antibodies lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added.
Antibody Type	Secondary Antibody
Application Notes	ELISA,Immunocytochemistry,Immunohistochemistry (frozen),(In)direct immunofluorescence.