

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Human lactoferrin, Clone: [Polyclonal], Monoclonal NMB-GAHU/LFR

Article Name	Goat anti Human lactoferrin, Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-GAHU/LFR
Supplier Catalog Number	GAHu/Lfr
Alternative Catalog Number	NMB-GAHU/LFR
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Product Description	In immunoelectrophoresis against human milk a single precipitin line is obtained. The antiserum does not react with any other protein component of human serum or plasma. In precipitating techniques as immunoelectrophoresis, single and double radial i...
Clonality	Monoclonal
Clone Designation	[Polyclonal]
UniProt	P02788
Buffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized ant
Source	Exocrine organs produce various secretions, each with its characteristic function. Proteins found in secretions may be divided into two groups: those specific for the particular secretion, and plasma proteins independent of the type of exocrine cells. Lac

Formula	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added.
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 µl human serum against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter center well and 10 µl of the antiserum in 3 mm diameter peripheral wells. In single radial immunodiffusion use 1% antiserum in the gel.