

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Human prealbumin, Clone: [Polyclonal], Monoclonal NMB-GAHU/PALB

| | |
|----------------------------|--|
| Article Name | Goat anti Human prealbumin, Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-GAHU/PALB |
| Supplier Catalog Number | GAHu/pAlb |
| Alternative Catalog Number | NMB-GAHU/PALB |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Product Description | The defined antibody specificity is directed to transthyretin as tested against human sera. In immunoelectrophoresis and double radial immunodiffusion (Ouchterlony), using various antiserum concentrations against appropriate concentrations of the imm... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| UniProt | P02766 |
| Buffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those in pooled goat serum. No foreign proteins added. Reconstitute the lyophilized antiseru |
| Source | Transthyretin is the fastest protein in the immunoelectrophoresis pattern of human serum. It consists of four identical polypeptide chains with no carbohydrate The molecular weight is 55,000 and the concentration in normal serum ranges from 0.1 to 0.4 mg |

| | |
|-------------------|--|
| Formula | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those in pooled goat serum. No foreign proteins added. |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 µl serum, plasma or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samp |