

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Human Secretory Component, free and bound-FITC, MinX none NMB-GAHU/SC/FITC

| Article Name | Goat IgG anti-Human Secretory Component, free and bound-FITC, MinX none |
|----------------------------|--|
| Biozol Catalog Number | NMB-GAHU/SC/FITC |
| Supplier Catalog Number | GAHu/SC/FITC |
| Alternative Catalog Number | NMB-GAHU/SC/FITC |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Species Reactivity | Human |
| Conjugation | FITC |
| Format | IgG |
| Target Specificity | Secretory Component - free and bound determinants |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | Tested in immunoelectrophoresis, double radial immunodiffusion and ELISA against a panel of appropriate secretions and purified Ig isotypes. The antiserum reacts with both bound secretory component (secretory IgA) and with the free SC present in huma |
| Clonality | Polyclonal |
| Clone Designation | [Polyclonal] |

| Buffer | Fluorochrome-coupled purified hyperimmune IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2) No preservative added, as it may interfere with the antibody activity. |
|-------------------|--|
| Source | Secretory component is present in human secretions bound to secretory IgA (slgA) and in free form. Secretory IgA (slgA) functions as a dimer or polymer and accounts for almost all specific mucosal antibody activity. A molecule of slgA is made up of two m |
| Formula | Fluorochrome-coupled purified hyperimmune IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2) No preservative added, as it may interfere with the antibody activity. |
| Antibody Type | Secondary Antibody |
| Application Notes | ELISA,Immunocytochemistry,Immunohistochemistry (frozen),(In)direct immunofluorescence. The lyophilized conjugate is shipped at ambient temperature and may be stored at +4C, prolonged storage at or below -20C. It is reconstituted by adding 1 ml sterile |