

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Human thyroxine binding globulin, Clone: [Polyclonal], Monoclonal NMB-GAHU/TBG

Article Name	Goat anti Human thyroxine binding globulin, Clone: [Polyclonal],
Allicie Name	Monoclonal
Biozol Catalog Number	NMB-GAHU/TBG
Supplier Catalog Number	GAHu/TBG
Alternative Catalog Number	NMB-GAHU/TBG
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Product Description	The reactivity of the antiserum is restricted to thyroxine binding globulin. In immunoelectrophoresis and radial immunodiffusion (Ouchterlony), using various antiserum concentrations against fresh normal human plasma a single precipitin line is obtai
Clonality	Monoclonal
Clone Designation	[Polyclonal]
UniProt	P05543
Buffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized an

Source	Thyroxine binding is a protein with a molecular weight of 55,700 kD. The protein is binding thyroxine as well as triiodothyronine. In immunoelectrophoresis it has a alpha mobility. The mean concentration in normal human serum is about 40 μ g/ml. Elevated
Formula	Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added.
Application Notes	Precipitation assays. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affec